

**A POTENTIAL INFRARED PROBLEM WITH THE DAMPING RATES
FOR GLUONS WITH SOFT MOMENTUM IN HOT QCD**

A. Abada, O. Azi and K. Benchallal

*Département de Physique, École Normale Supérieure,
BP 92, Vieux Kouba 16050, Alger, Algeria*

enskppps@ist.cerist.dz

Abstract

We calculate the damping rate γ_l for longitudinal gluons with zero momentum in finite high temperature QCD and show that some of its contributing terms are infrared divergent. This is in contrast with the expectation that this damping rate is to be equal to the corresponding one γ_t for transverse gluons which is known to be finite. Our calculation was motivated by the fact that similar divergent terms occur when we calculated in a previous work γ_t to order p^2 , $\mathbf{p}$ being the momentum of the gluon. After we present our results, we briefly discuss them.

In a previous work [1], we undertook the calculation of the damping rate $\gamma_t(p)$ for transverse gluons with soft momentum $\mathbf{p}$ in finite-high-temperature QCD. Knowing that γ_t at zero momentum is finite [2], our main aim in that work was to argue that for small enough $p = |\mathbf{p}|$, the expansion of $\gamma_t(p)$ in powers of $(p/m_g)^2$ is valid, m_g being the inverse gluonic correlation length. We suggested that, when contrasted with a previously recent calculation [3] based on letting p come from the hard limit (set by the temperature T) down towards the interior of the soft region (set by $m_g \sim gT$ where g is the QCD coupling constant), a calculation that produced a behavior of the form $\ln(1/g)$ for $\gamma_t(p)$, such a result could possibly be indicative of a kind of transition around some ‘critical’ scale. We therefore expanded $\gamma_t(p)$ in the following manner:

$$\gamma_t(p) = \frac{g^2 N_c T}{24\pi} \left[a_{t0} + a_{t1} \left(\frac{p}{m_g} \right)^2 + \dots \right], \quad (1)$$

and gave an analytic expression for a_{t1} (the first coefficient a_{t0} was already calculated in [2]). N_c is the number of colors.

The natural step forward was to calculate numerically a_{t1} . But quite interestingly, when we started manipulating the expression of a_{t1} we obtained, we realized that some of its contributing terms are infrared divergent (in a sense that will become clear later in the text). Thus, on the one hand, the quick conclusion is that a_{t1} may be infinite and hence, the expansion in (1) is not valid after all. On the other hand, such a potential infrared divergence seems in contradiction with the physical expectation that the quark-gluon plasma is stable for, at least, very small gluonic momenta.

In order to look into the matter more closely, we undertake in this work the derivation of $\gamma_l(0)$, the damping rate for longitudinal gluons at zero momentum. Our motivation is that, first, this damping rate is to be equal to $\gamma_t(0)$, simply because at zero momentum, there is no difference between transverse and longitudinal directions. Second, the interesting point in this calculation is that in order to get γ_l at zero p , it is already necessary to expand it to order $(p/m_g)^2$. After the calculation is performed, we indeed recover the finite analytic expression that is equal to $\gamma_t(0)$, but, surprisingly enough, together with additional terms quite similar in form to the ones we obtained in the case of $\gamma_t(p)$, terms when taken individually present an infrared divergence. To the best of our knowledge, in this context, such infrared-divergent terms have not been discussed previously in the literature. Before furthering this discussion, we first present in the sequel our results.

In the imaginary-time formalism, the euclidian momentum of the gluon is $P^\mu = (p_0, \mathbf{p})$ such that $P^2 = (p_0)^2 + p^2$ with $\mathbf{p} = p \hat{\mathbf{p}}$ and $p_0 = 2\pi nT$ where n is an integer. Real-time amplitudes are obtained via the analytic continuation $p_0 = -i\omega + 0^+$ where ω is the energy of the gluon. A momentum is said to be soft if both ω and p are of order gT ; it is said to be hard if one is or both are of order T . In the Coulomb gauge, the complete inverse gluon propagator is:

$$D_{\mu\nu}^{-1}(P) = P^2 \delta^{\mu\nu} - P^\mu P^\nu - \Pi^{\mu\nu}(P) + \frac{1}{\xi_C} \delta^{\mu i} \delta^{\nu j} p^i p^j, \quad (2)$$

where $\Pi^{\mu\nu}(P)$ is the gluon self-energy and the last term is due to Coulomb-gauge fixing. We work in the

strict Coulomb gauge $\xi_C = 0$. The gluon self-energy can be decomposed into:

$$\Pi^{\mu\nu}(P) = \delta\Pi^{\mu\nu}(P) + {}^*\Pi^{\mu\nu}(P) , \quad (3)$$

where $\delta\Pi$ is the hard-thermal-loop and ${}^*\Pi$ is the effective self-energy [4]. P being soft, the hard thermal loop is of the same order of magnitude as the inverse free propagator, i.e., $\delta\Pi \sim (gT)^2$, while the effective self-energy is of an order of magnitude higher, i.e., ${}^*\Pi \sim g(gT)^2$. Since the momentum running inside ${}^*\Pi$ is soft, to calculate it, we have to use effective vertices and propagators instead of their bare (tree) counterparts. This ensures the correct expression for the $g(gT)^2$ -correction to the inverse gluon propagator and, in particular, that this correction is gauge-independent [4].

The hard-thermal-loop $\delta\Pi$ is already known from the literature [2,4]. It is real and contributes to the determination of the spectrum of the soft gluonic excitations to leading order gT . More explicitly, $\delta\Pi$ is gauge-invariant and can be expressed in terms of only two independent scalar functions, i.e.,:

$$\begin{aligned} \delta\Pi^{00}(P) &= \delta\Pi_l(P) ; & \delta\Pi^{0i}(P) &= -\frac{p_0 p^i}{p^2} \delta\Pi_l(P) ; \\ \delta\Pi^{ij}(P) &= (\delta^{ij} - \hat{p}^i \hat{p}^j) \delta\Pi_t(P) + \hat{p}^i \hat{p}^j \frac{(p_0)^2}{p^2} \delta\Pi_l(P) . \end{aligned} \quad (4)$$

The expressions of $\delta\Pi_l(P)$ and $\delta\Pi_t(P)$ read [2]:

$$\delta\Pi_l(P) = 3m_g^2 Q_1\left(\frac{ip_0}{p}\right) ; \quad \delta\Pi_t(P) = \frac{3}{5}m_g^2 \left[Q_3\left(\frac{ip_0}{p}\right) - Q_1\left(\frac{ip_0}{p}\right) - \frac{5}{3} \right] , \quad (5)$$

where the Q_n are Legendre functions of the second kind. The thermal gluonic mass is given explicitly by $m_g = \sqrt{N_c + (1/2)N_f} gT/3$, where N_f is the number of flavors.

The effective propagator for soft gluons that intervenes in the calculation of the effective self-energy is obtained by inverting (2) while disregarding ${}^*\Pi$. In the strict Coulomb gauge, its nonzero components are ${}^*\Delta_C^{00}(P) = {}^*\Delta_l(P)$ and ${}^*\Delta_C^{ij}(P) = (\delta^{ij} - \hat{p}^i \hat{p}^j) {}^*\Delta_t(P)$, where ${}^*\Delta_t$ and ${}^*\Delta_l$ are given by:

$${}^*\Delta_t(P) = \frac{1}{P^2 - \delta\Pi_t(P)} ; \quad {}^*\Delta_l(P) = \frac{1}{p^2 - \delta\Pi_l(P)} . \quad (6)$$

After analytic continuation to real energies, the pole in ω of ${}^*\Delta_{t(l)}$ yields the dispersion relation $\omega_{t(l)}(p)$ of the transverse (longitudinal) gluons to order gT . The longitudinal dispersion equation can be rewritten in the following explicit manner:

$$\ln \frac{\omega_l + p}{\omega_l - p} = \frac{2p}{3\omega_l} \left(3 + \frac{p^2}{m_g^2} \right) . \quad (7)$$

One finds for soft longitudinal gluons:

$$\omega_l(p) = m_g \left[1 + \frac{3}{10} \left(\frac{p}{m_g} \right)^2 - \frac{3}{280} \left(\frac{p}{m_g} \right)^4 + \frac{1}{6000} \left(\frac{p}{m_g} \right)^6 + \frac{489}{43120000} \left(\frac{p}{m_g} \right)^8 + \dots \right] . \quad (8)$$

To order gT , the gluons are not damped. The leading order of the damping rates is obtained after is included in the dispersion equations the contribution from the effective self-energy ${}^*\Pi$, which of course has a

more complicated structure than that of $\delta\Pi$. One shows that in the strict Coulomb gauge, it also depends on two scalar functions such that the mass-shell condition that determines the longitudinal dispersion relation is:

$$p^2 - \delta\Pi_l(-i\Omega_l, p) - {}^*\Pi_l(-i\Omega_l, p) = 0 , \quad (9)$$

where ${}^*\Pi_l = {}^*\Pi^{00}$. The longitudinal damping rate is defined by $\gamma_l(p) \equiv -\text{Im } \Omega_l(p)$. Since it is g -times smaller than the energy $\omega_l(p)$, we can write from (9):

$$\gamma_l(p) = \frac{\text{Im} {}^*\Pi_l(-i\omega, p)}{\frac{\partial}{\partial\omega} \delta\Pi_l(-i\omega, p)} \Big|_{\omega=\omega_l(p)+i0^+} . \quad (10)$$

The denominator in (9) is easy to get from (5). One has $\partial/\partial\omega \delta\Pi(-i\omega, p)|_{\omega=\omega_l(p)} = -2m_g(p/m_g)^2 + \dots$. This means that in order to get $\gamma_l(p=0)$, we have to expand the imaginary part of the longitudinal self-energy to order p^2 .

In the Coulomb gauge, the only diagrams that contribute to $\text{Im} {}^*\Pi_l(P)$ above the light cone are the three-gluon (3g) and four-gluon (4g) one loop-diagrams with soft internal momentum [4]. Hence we write:

$$\begin{aligned} \text{Im } {}^*\Pi_l(P) = & -\frac{g^2 N_c}{2} \text{Im Tr}_{\text{soft}} \left[{}^*\Gamma^{00\lambda\sigma}(P, -P, K, -K) {}^*\Delta_C^{\lambda\sigma}(K) \right. \\ & \left. + {}^*\Gamma^{\sigma 0\lambda}(-Q, P, -K) {}^*\Delta_C^{\lambda\lambda'}(K) {}^*\Gamma^{\lambda'0\sigma'}(-K, P, -Q) {}^*\Delta_C^{\sigma'\sigma}(Q) \right] , \end{aligned} \quad (11)$$

where K is the internal loop-momentum, $Q = P - K$ and $\text{Tr} \equiv T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3}$. The subscript ‘soft’ means that only soft values of k are allowed in the integral. Eq (11) is what one would normally write for the 3g and 4g contributions to the imaginary part of the gluon self-energy, except that everywhere, tree quantities are replaced by the corresponding effective ones. The gluonic effective vertices can be written in the following form:

$${}^*\Gamma^{(n)} = \Gamma^{(n)} + \delta\Gamma^{(n)} ; \quad n = 3, 4 , \quad (12)$$

where the first term is the tree QCD gluon vertex and the second one sums up the contributions from hard thermal loops with n external legs. In the case $n = 3$ it can be written as:

$$\delta\Gamma^{\mu\nu\lambda}(-Q, P, -K) = -\delta\Gamma^{\mu\nu\lambda}(-K, P, -Q) = 3m_g^2 \int \frac{d\Omega_S}{4\pi} \frac{S^\mu S^\nu S^\lambda}{PS} \left(\frac{iq_0}{QS} - \frac{ik_0}{KS} \right) , \quad (13)$$

where $S \equiv (i, \hat{\mathbf{s}})$ and Ω_S is the solid angle of the unit three-vector $\hat{\mathbf{s}}$. Also, $PS = ip_0 + \mathbf{p} \cdot \hat{\mathbf{s}}$, etc. In the case $n = 4$ we have:

$$\delta\Gamma^{\mu\nu\lambda\sigma}(P, -P, K, -K) = 3m_g^2 \int \frac{d\Omega_S}{4\pi} \frac{S^\mu S^\nu S^\lambda S^\sigma}{PS KS} \left(\frac{ip_0 - ik_0}{PS - KS} - \frac{ip_0 + ik_0}{PS + KS} \right) . \quad (14)$$

To be complete, we give the expression of the 3g tree vertex:

$$\begin{aligned} \Gamma^{\mu\nu\lambda}(-Q, P, -K) &= -\Gamma^{\lambda\nu\mu}(-K, P, -Q) \\ &= (P + K)^\mu \delta^{\nu\lambda} + (Q - K)^\nu \delta^{\lambda\mu} - (P + Q)^\lambda \delta^{\mu\nu} , \end{aligned} \quad (15)$$

and that of the $4g$ tree vertex:

$$\Gamma^{\mu\nu\lambda\sigma}(P, -P, K, -K) = 2\delta^{\mu\nu}\delta^{\lambda\sigma} - \delta^{\mu\lambda}\delta^{\nu\sigma} - \delta^{\mu\sigma}\delta^{\nu\lambda}. \quad (16)$$

From eq (11) and the discussion before eq (6), we have:

$$\begin{aligned} \text{Im } {}^*\Pi_l(P) = & -\frac{g^2 N_c}{2} \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} \left[{}^*\Gamma^{0000}(P, -P, K, -K) {}^*\Delta_l(K) \right. \\ & + {}^*\Gamma^{00ij}(P, -P, K, -K) (\delta^{ij} - \hat{k}^i \hat{k}^j) {}^*\Delta_t(K) \\ & + {}^*\Gamma^{000}(-Q, P, -K) {}^*\Delta_l(K) {}^*\Gamma^{000}(-K, P, -Q) {}^*\Delta_l(Q) \\ & + {}^*\Gamma^{00i}(-Q, P, -K) (\delta^{ij} - \hat{k}^i \hat{k}^j) {}^*\Delta_t(K) {}^*\Gamma^{j00}(-K, P, -Q) {}^*\Delta_l(Q) \\ & + {}^*\Gamma^{j00}(-Q, P, -K) {}^*\Delta_l(K) {}^*\Gamma^{00i}(-K, P, -Q) (\delta^{ij} - \hat{q}^i \hat{q}^j) {}^*\Delta_t(Q) \\ & \left. + {}^*\Gamma^{n0i}(-Q, P, -K) (\delta^{ij} - \hat{k}^i \hat{k}^j) {}^*\Delta_t(K) {}^*\Gamma^{j0m}(-K, P, -Q) (\delta^{mn} - \hat{q}^m \hat{q}^n) {}^*\Delta_t(Q) \right]. \quad (17) \end{aligned}$$

There are six contributions: two from the $4g$ vertices and four from the $3g$ vertices. We will take henceforth $m_g \equiv 1$ and all momenta and energies are in units of it. This simplifies considerably the expressions we work with and, if and when needed, the m_g -dependence can easily be recovered in the final results. Each contribution has to be calculated separately. As an illustration, let us discuss how we manipulate the contribution from the $3g$ vertices in which the two effective propagators involved are both longitudinal. The corresponding term in (17) is:

$$\text{Im } {}^*\Pi_{l3gll}(P) = -\frac{g^2 N_c}{2} \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} {}^*\Gamma^{000}(-Q, P, -K) {}^*\Delta_l(K) {}^*\Gamma^{000}(-K, P, -Q) {}^*\Delta_l(Q), \quad (18)$$

in a clear notation. Using eqs (12), (13) and (15), we have:

$$\begin{aligned} \text{Im } {}^*\Pi_{l3gll}(P) = & -\frac{9}{2} g^2 N_c \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} {}^*\Delta_l(K) {}^*\Delta_l(Q) \\ & \times \left[\left(\int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} \right)^2 - \int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} \int \frac{d\Omega_S}{4\pi} \frac{iq_0}{PSQS} \right]. \quad (19) \end{aligned}$$

We need an expression for the two solid-angle integrals involved in (19). For this purpose, we use the simple expansion $1/PS = 1/ip_0 (1 - \mathbf{p} \cdot \hat{\mathbf{s}}/ip_0 - \mathbf{p} \cdot \hat{\mathbf{s}}^2/p_0^2 + \dots)$, which is valid as long as $p < |ip_0|$, a condition satisfied in the region $p < m_g$ before analytic continuation to real energies and after. We then perform the angular integrals straightforwardly and get:

$$\begin{aligned} \int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} = & \frac{ik_0}{ip_0 k} \left[Q_{0k} - \frac{px}{ip_0} \left(1 - \frac{ik_0}{k} Q_{0k} \right) \right. \\ & \left. - \frac{p^2}{p_0^2} \left[-x^2 \frac{ik_0}{k} \left(1 - \frac{ik_0}{k} Q_{0k} \right) + \frac{1}{2} (1 - x^2) \left(\frac{ik_0}{k} + \left(1 + \frac{k_0^2}{k^2} \right) Q_{0k} \right) \right] + \dots \right], \quad (20) \end{aligned}$$

where Q_{0k} stands for $Q_0(\frac{ik_0}{k})$ and $x = \cos(\hat{\mathbf{p}}, \hat{\mathbf{k}})$. We obtain a similar expression for the other solid-angle integral: we only need to replace K by Q .

The next step is to multiply the expressions for the solid-angle integrals as needed in eq (19) while still expanding in powers of p . Then we integrate over the solid angle of $\hat{\mathbf{k}}$ and the odd-powered terms in p cancel out. For the first piece in (19) we get:

$$\begin{aligned}
& -9 g^2 N_c \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} {}^* \Delta_l(K) {}^* \Delta_l(Q) \left(\int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} \right)^2 = \frac{9 g^2 N_c}{2\pi^2} \text{Im } T \sum_{k_0} \int_0^\infty \frac{k^2}{-p_0^2} dk \\
& \quad \times {}^* \Delta_l(K) \left[\frac{k_0^2}{k^2} Q_{0k}^2 + \frac{p^2}{3} \left[\frac{k_0^2}{k^2} Q_{0k}^2 \left(\frac{1}{k} \partial_k + \frac{1}{2} \partial_k^2 \right) + \frac{2}{ip_0} \frac{k_0^2}{k^2} Q_{0k} \left(1 - \frac{ik_0}{k} Q_{0k} \right) \partial_k \right. \right. \\
& \quad - \frac{1}{p_0^2} \left(\frac{k_0^2}{k^2} \left(1 - \frac{ik_0}{k} Q_{0k} \right)^2 - 2 \frac{ik_0^3}{k^3} Q_{0k} \left(1 - \frac{ik_0}{k} Q_{0k} \right) \right. \\
& \quad \left. \left. + 2 \frac{k_0^2}{k^2} Q_{0k} \left(\frac{ik_0}{k} + \left(1 + \frac{k_0^2}{k^2} \right) Q_{0k} \right) \right] + \dots \right] {}^* \Delta_l(q_0, k), \tag{21}
\end{aligned}$$

where ∂_k stands for $\partial/\partial k$. Similarly, the second piece in (19) reads:

$$\begin{aligned}
& +9 g^2 N_c \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} {}^* \Delta_l(K) {}^* \Delta_l(Q) \int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} \int \frac{d\Omega_S}{4\pi} \frac{iq_0}{PSQS} \\
& = \frac{9 g^2 N_c}{2\pi^2} \text{Im } T \sum_{k_0} \int \frac{k^2}{-p_0^2} dk {}^* \Delta_l(K) \left[\left(1 + \frac{k^2}{3} \right)^2 + \frac{p^2}{3} \left[\left(1 + \frac{k^2}{3} \right) \left(1 + \frac{iq_0}{ip_0} \right) \right. \right. \\
& \quad - \frac{2k^2}{9} \frac{ik_0}{ip_0} - \frac{2}{p_0^2} \left(1 + \frac{k^2}{3} \right)^2 + \frac{ik_0 iq_0 k^2}{9 p_0^2} + \left(\left(1 + \frac{k^2}{3} \right)^2 \right. \\
& \quad \left. \left. + \frac{k^2}{3} \left(2 - \frac{ik_0}{ip_0} + \frac{iq_0}{ip_0} \right) \left(1 + \frac{k^2}{3} \right) \right) \frac{1}{k} \partial_k + \frac{1}{2} \left(1 + \frac{k^2}{3} \right)^2 \partial_k^2 \right] + \dots \left] {}^* \Delta_l(q_0, k). \tag{22}
\end{aligned}$$

In order to eliminate the appearance of Q_{0k}^2 in (21) [1], extra work is needed. For this purpose, we use the expression for ${}^* \Delta_l(K)^{-1}$ one obtains using eqs (6) and (5). After some algebra and disregarding terms that from the outset do not yield an imaginary part to the effective self-energy after analytic continuation, we get:

$$\begin{aligned}
& -9 g^2 N_c \text{Im } T \sum_{k_0} \int \frac{d^3 k}{(2\pi)^3} {}^* \Delta_l(K) {}^* \Delta_l(Q) \left(\int \frac{d\Omega_S}{4\pi} \frac{ik_0}{PSKS} \right)^2 \\
& = \frac{9 g^2 N_c}{2\pi^2} \text{Im } T \sum_{k_0} \int \frac{k^2}{-p_0^2} dk \left[- \left(1 + \frac{k^2}{3} \right)^2 {}^* \Delta_l(K) + \frac{ik_0}{3k} Q_{0k} \right. \\
& \quad + \frac{p^2}{3} \left[\frac{1}{p_0^2} \text{Big} \left(2 + \frac{4k^2}{3} + \frac{2k^4}{9} - \frac{k_0^2 k^2}{9} \right) {}^* \Delta_l(K) + \frac{1}{3} \left(-2 + \frac{k_0^2}{k^2} \right) \frac{ik_0}{k} Q_{0k} \right. \\
& \quad + \left(- \left(1 + \frac{k^2}{3} \right) \left(1 + \frac{k^2}{3} \left(1 - 2 \frac{ik_0}{ip_0} \right) \right) {}^* \Delta_l(K) + \frac{1}{3} \left(1 - 2 \frac{ik_0}{ip_0} \right) \frac{ik_0}{k} Q_{0k} \right) \frac{1}{k} \partial_k \\
& \quad \left. \left. + \frac{1}{2} \left(- \left(1 + \frac{k^2}{3} \right)^2 {}^* \Delta_l(K) + \frac{ik_0}{3k} Q_{0k} \right) \partial_k^2 \right] + \dots \right] {}^* \Delta_l(q_0, k). \tag{23}
\end{aligned}$$

Remember that all momenta and energies are in units of m_g , set to one. Putting the two pieces (22) and (23) together, we find:

$$\begin{aligned}
& \text{Im } {}^* \Pi_{l3gl}(P) = \frac{9 g^2 N_c}{2\pi^2} \text{Im } T \sum_{k_0} \int \frac{k^2}{-p_0^2} dk \left[\frac{ik_0}{3k} Q_{0k} + \frac{p^2}{3} \left[\left(\left(1 + \frac{k^2}{3} \right) \left(1 + \frac{iq_0}{ip_0} \right) - \frac{ik_0 k^2}{3 ip_0} \right) {}^* \Delta_l(K) \right. \right. \\
& \quad + \frac{1}{3 p_0^2} \left(-2 + \frac{k_0^2}{k^2} \right) \frac{ik_0}{k} Q_{0k} + k^2 \left(1 + \frac{k^2}{3} \right) {}^* \Delta_l(K) \partial_k + \frac{1}{3k} \left(1 - 2 \frac{ik_0}{ip_0} \right) \frac{ik_0}{k} Q_{0k} \partial_k \\
& \quad \left. \left. + \frac{ik_0}{6k} Q_{0k} \partial_k^2 \right] \dots \right] {}^* \Delta_l(q_0, k). \tag{24}
\end{aligned}$$

The next step is to perform the sum over k_0 . We do this by using the spectral representations [5] of the different quantities involved. In particular, we have for the effective propagators:

$$^*\Delta_{t,l}(K) = \int_0^{\frac{1}{T}} d\tau e^{ik_0\tau} \int_{-\infty}^{+\infty} d\omega \rho_{t,l}(\omega, k) (1 + n(\omega)) e^{-\omega\tau}, \quad (25)$$

where $n(\omega) = 1/(\exp(\omega/T) - 1)$ is the Bose-Einstein distribution and the spectral densities $\rho_{t,l}(\omega, k)$ have the following form:

$$\rho_{t,l}(\omega, k) = \frac{Z_{t,l}(k)}{2\omega_{t,l}(k)} \left[\delta(\omega - \omega_{t,l}(k)) - \delta(\omega + \omega_{t,l}(k)) \right] + \beta_{t,l}(\omega, k) \Theta(k^2 - \omega^2), \quad (26)$$

where Θ is the step function. The two residues $Z_{t,l}(k)$ and the two cuts $\beta_{t,l}(\omega, k)$ are given in eqs (3.23) and (3.24) of [1]; see also [5]. The sum over k_0 can now be performed straightforwardly. The analytic continuation to real energies is obtained by the subsequent replacement $ip_0 \rightarrow \omega_l(p) + i0^+$ and we extract the imaginary part from our expressions using the known relation $1/(x + i0^+) = \text{Pr}(1/x) - i\pi\delta(x)$. We find:

$$\begin{aligned} \text{Im } ^*\Pi_{l3gl}(P) = & \frac{3g^2 N_c T}{4\pi} \int_0^{+\infty} dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(\omega_l - \omega_1 - \omega_2) \left[-\omega_l \omega_1 k \Theta_1 \right. \\ & - p^2 \left[-k^2 \left(3 + \frac{2}{3} k^2 \right) \rho_{l1} + \frac{\omega_1 k}{3} \left(2 + \frac{\omega_1^2}{k^2} \right) \Theta_1 - 2k^3 \left(1 + \frac{k^2}{3} \right) \rho_{l1} \partial_k \right. \\ & \left. \left. + \frac{\omega_1}{3} (1 - 2\omega_1) \Theta_1 \partial_k + \frac{\omega_1 k}{6} \Theta_1 \partial_k^2 \right] + \dots \right] \rho_{l2}. \end{aligned} \quad (27)$$

In this equation, ρ_{li} stands for $\rho_l(\omega_i, k)$, $i = 1, 2$, and $\Theta_1 = \Theta(k^2 - \omega_1^2)$. Also, for ω soft, we have used $n(\omega) \simeq T/\omega$. Finally, note that ω_l depends on p and in principle eq (27) still needs a further expansion. But this will prove unnecessary.

The other contributions to $\text{Im } ^*\Pi_l(P)$ are calculated in a similar manner. The expressions of the two $4g$ -contributions are quicker to obtain whereas more work is needed for the other $3g$ -contributions, especially the $3gtt$ -part. We spare the reader the details and here we give the final expressions. The two $4g$ -contributions read:

$$\begin{aligned} \text{Im } ^*\Pi_{l4gl}(P) = & \frac{3g^2 N_c T}{4\pi} \int_0^{\infty} dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(\omega_l - \omega_1 - \omega_2) \left[\omega_l \omega_1 k \Theta_1 \right. \\ & \left. - p^2 \left[-\omega_1 k \Theta_1 + \frac{4}{3} \omega_1 k^2 \epsilon(\omega_1) \delta_1 - \frac{2}{3} \omega_1^2 k^2 \epsilon(\omega_1) \partial_{\omega_1^2} \delta_1 \right] + \dots \right] \rho_{l2}; \end{aligned} \quad (28)$$

$$\begin{aligned} \text{Im } ^*\Pi_{l4gt}(P) = & \frac{3g^2 N_c T}{4\pi} \int_0^{\infty} dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(\omega_l - \omega_1 - \omega_2) \left[-\omega_l \omega_1 k \left(1 - \frac{\omega_1^2}{k^2} \right) \Theta_1 \right. \\ & \left. - p^2 \left[\frac{\omega_1}{k^3} \left(k^4 - \frac{k^2}{3} + \frac{4}{3} k^2 \omega_1 - k^2 \omega_1^2 \right) \Theta_1 + \frac{2}{3} \omega_1^2 \epsilon(\omega_1) \delta_1 \right] + \dots \right] \rho_{l2}, \end{aligned} \quad (29)$$

Where $\epsilon(\omega)$ is the sign function and $\delta_1 \equiv \delta(\omega_1^2 - k^2)$. As to the $3g$ -contributions, we have:

$$\begin{aligned} \text{Im } ^*\Pi_{l3gtl}(P) = \text{Im } ^*\Pi_{l3gl}(P) = & \frac{g^2 N_c T}{24\pi} \int_0^{\infty} dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(1 - \omega_1 - \omega_2) \\ & \times p^2 \left[9k^2 (k^2 - \omega_1^2)^2 \rho_{l1} \rho_{l2} - \frac{3\omega_1}{k} (k^2 - \omega_1^2) \Theta_1 \rho_{l2} + \frac{3\omega_1}{2k^3} (k^2 + 2\omega_1 - \omega_1^2)^2 \Theta_1 \rho_{l2} \right] + \dots, \end{aligned} \quad (30)$$

and we have:

$$\begin{aligned}
\text{Im } \Pi_{l3gtt}(P) = & \frac{3g^2 N_c T}{4\pi} \int_0^\infty dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(\omega_l - \omega_1 - \omega_2) \left[\omega_l \omega_1 k \left(1 - \frac{\omega_1^2}{k^2}\right) \Theta_1 \right. \\
& - p^2 \left[\frac{2}{9} (3k^2(1 - 2k^2) + (1 + 6k^2)\omega_1\omega_2 + 8\omega_1^2\omega_2^2) \rho_{t1} + \frac{\omega_1}{3k^3} (k^2 - 2k^4 - 2k^2\omega_1 \right. \\
& + (k^2 - 1)\omega_1^2 + 2\omega_1^3 + \omega_1^4) \Theta_1 + \frac{2k}{9} (1 - 2k^2 - 6k^4 - 2\omega_1 + 2(1 + 4k^2)\omega_1^2 - 2\omega_1^4) \rho_{t1} \partial_k \\
& + \frac{\omega_1}{3} \left(-1 + 2\omega_1 + \frac{\omega_1}{k^2} - \frac{2\omega_1^2}{k^2} \right) \Theta_1 \partial_k + \frac{k^2}{9} (1 - 2k^2 + 4(-1 + k^2)\omega_1 + 6\omega_1^2 - 4\omega_1^3) \rho_{t1} \partial_k^2 \\
& \left. \left. - \frac{\omega_1 k}{6} \left(1 - \frac{\omega_1^2}{k^2}\right) \Theta_1 \partial_k^2 \right] + \dots \right] \rho_{t2} .
\end{aligned} \tag{31}$$

We put now the different contributions together. The terms independent of p cancel out as they should and, after reorganizing the terms proportional to p^2 in a suitable manner, we obtain, using eq (10) in the limit $p \rightarrow 0$:

$$\gamma_l(0) = \frac{g^2 N_c T}{24\pi} \left[a_{l0}^{(1)} + a_{l0}^{(2)} \right] , \tag{32}$$

where the pure numbers $a_{l0}^{(1)}$ and $a_{l0}^{(2)}$ have the following analytic expressions:

$$\begin{aligned}
a_{l0}^{(1)} = & 9 \int_0^\infty dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(1 - \omega_1 - \omega_2) \left[k^4 \rho_{l1} \rho_{l2} \right. \\
& \left. - k^2 (k^2 - \omega_1^2)^2 \rho_{t1} \rho_{l2} + 2(k^2 + \omega_1\omega_2)^2 \rho_{t1} \rho_{t2} + \frac{\omega_1}{6k^3} (k^2 - \omega_1^2)^2 \Theta_1 \rho_{t2} \right] ,
\end{aligned} \tag{33}$$

and:

$$\begin{aligned}
a_{l0}^{(2)} = & 9 \int_0^\infty dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(1 - \omega_1 - \omega_2) \left[\frac{\omega_1}{3} (1 - 2\omega_1) \Theta_1 \partial_k \rho_{l2} + \frac{\omega_1 k}{6} \Theta_1 \partial_k^2 \rho_{l2} \right. \\
& + \frac{\omega_1}{3} \left(1 - 2\omega_1 - \frac{\omega_1^2}{k^2} + 2\frac{\omega_1^3}{k^2} \right) \partial_k \Theta_1 \rho_{t2} + \frac{\omega_1 k}{3} \left(1 - \frac{\omega_1^2}{k^2} \right) \partial_k \Theta_1 \partial_k \rho_{t2} + \frac{\omega_1}{6k} (k^2 - \omega_1^2) \partial_k^2 \Theta_1 \rho_{t2} \\
& + \partial_k \left[-\frac{2k^3}{3} \left(1 + \frac{k^2}{3} \right) (2 - \omega_1) \rho_{l1} \rho_{l2} + \frac{2k}{9} (k^2(1 - 3k^2) + (1 - 4k^2)\omega_1\omega_2 - \omega_1^2\omega_2^2) \rho_{t1} \rho_{t2} \right. \\
& + \frac{2\omega_1^2}{3} \left(2 + \frac{\omega_1}{k^2} - \frac{\omega_1^2}{k^2} \right) \Theta_1 \rho_{t2} + \frac{k^2}{9} (1 - 2k^2 - 4(1 - k^2)\omega_1 + 6\omega_1^2 - 4\omega_1^3) \rho_{t1} \partial_k \rho_{t2} \\
& \left. \left. - \frac{\omega_1}{6} \partial_k \left(\left(k - \frac{\omega_1}{k} \right) \Theta_1 \rho_{t2} \right) \right] + \frac{4}{3} k^2 \omega_1 \epsilon(\omega_1) \delta_1 \rho_{l2} - \frac{2}{3} k^2 \omega_1^2 \epsilon(\omega_1) \partial_{\omega_1^2} \delta_1 \rho_{l2} + \frac{2}{3} \omega_1^2 \epsilon(\omega_1) \delta_1 \rho_{t2} \right] .
\end{aligned} \tag{34}$$

Things are now well set up to pursue our initial discussion. When introducing this work, we have said that at zero momentum, the longitudinal-gluon damping rate is to be equal to the transverse-gluon one, i.e., $\gamma_l(0) = \gamma_t(0)$. If one calculates the transverse-gluon damping rate at zero momentum, one finds $\gamma_t(0) = \frac{g^2 N_c T}{24\pi} a_{t0}$, where $a_{t0} = a_{l0}^{(1)}$, the expression of which is given in (33) above, see [2]. Therefore, one automatically expects $a_{l0}^{(2)}$ to vanish. But this seems not trivial an issue at all. Indeed, take for example the term $\omega_1/3 (1 - 2\omega_1) \Theta_1 \partial_k \rho_{l2}$ in the analytic expression of $a_{l0}^{(2)}$ given in (34). It turns out that the contribution of this term to $a_{l0}^{(2)}$ and hence to $\gamma_l(0)$ is infrared divergent. More precisely, if one cuts the lower bound of the k -integral to a small momentum $\eta \ll 1$, then one straightforwardly works out the contribution of this

term to $a_{l0}^{(2)}$ and finds:

$$\begin{aligned} \int_{\eta}^{\infty} dk \int_{-\infty}^{+\infty} \frac{d\omega_1}{\omega_1} \int_{-\infty}^{+\infty} \frac{d\omega_2}{\omega_2} \delta(1 - \omega_1 - \omega_2) \frac{1}{3} \omega_1 (1 - 2\omega_1) \Theta_1 \partial_k \rho_{l2} = -\frac{1}{6\eta^2} - \frac{1}{20\eta} + \frac{1}{20} \\ - \frac{1}{2} \int_{1/2}^{+\infty} dk \frac{1-2k}{k} \left[\left(3 + k^2 + \frac{3(1-k)}{2k} \ln(2k-1) \right)^2 + \left(\frac{3\pi(1-k)}{2k} \right)^2 \right]^{-1} + O(\eta) . \end{aligned} \quad (35)$$

Note that the integral over k in the right-hand side of the above equation is finite. Thus one clearly sees that this term is infrared divergent. Other terms have a similar behavior.

Remember that we decided to carry out this work when we noticed that terms similar to the ones in (34) do appear in the analytic expression of a_{t1} in eq (1), terms which are infrared divergent too [1]. At this stage, it becomes clear that the appearance of such terms is linked to the expansion in powers of p^2 . Besides, their appearance in the expression of $\gamma_l(0)$ is puzzling to a certain degree because as we said, $\gamma_l(0)$ is expected to be divergence-free. It therefore may be that though such terms, when taken individually, are infrared divergent, when put together, the infrared pieces cancel out and we are left with a finite contribution, likely a vanishing one in the case of $a_{l0}^{(2)}$. But it should be clear that checking through this point is a delicate matter. In any case, it is beyond the scope of the present work.

If such a cancellation does occur for both $a_{l0}^{(2)}$ and a_{t1} , does it signal to a more basic pattern related to some aspects of the infrared behavior of hot QCD or is it a mere ‘coincidence’? On the contrary, if the infrared divergences persist, is it possible to devise a sort of finite-temperature infrared-renormalization scheme that would absorb them or is it that we will have to ‘live with’, so to speak? If such a scheme is possible, how consistent would it be with other calculations?

Still in the case where these divergences persist, we can also naturally ask whether they are linked in some way to the logarithmic behavior $\ln(1/g)$ discussed in [3]; in other words, whether they are merely a different manifestation of the same phenomenon. We think it is too early to tackle thoroughly this interesting question but nevertheless, we can put forward the following argument. The logarithmic behavior is obtained when letting p come from above m_g down into the interior of the soft region, but keeping it larger than what is expected to be the magnetic scale, i.e., $g^2 T$. Therefore, though arguably we may think of the infrared divergences in a_{t1} as being another expression of this logarithmic behavior, this cannot be argued for $a_{l0}^{(2)}$ simply because here p has to be strictly zero and the logarithmic behavior cannot be extrapolated down to zero momentum. Besides, Since the infrared divergences we obtain in the case of $\gamma_l(0)$ are closer in nature to the ones we have for $\gamma_t(p)$, we find it difficult at this stage to answer this question positively, especially that the rôle of the magnetic effects is still somewhat obscure. In any case, one should not forget that the quark-gluon plasma is expected to be stable for zero and very small gluonic momenta, and therefore, any divergence in the gluonic damping rates, if it persists after scrutiny, has to be addressed from this standpoint.

It seems to us that all the issues we are raising in this work are worth investigating, especially that as far as we can tell, they haven’t been (at least fully) addressed in such a way previously.

We would like to thank Asmaa Abada and Olivier Pène from Orsay for interesting and quite useful exchanges that actually helped reshape parts of our discussion.

References

- [1] A. Abada, O. Azi, and A. Tadj, ‘*Damping rate for transverse gluons with finite soft momentum in hot QCD*’, ENSK-ph/97-02, [hep-ph/9710549](#), submitted to JHEP.
- [2] E. Braaten and R.D. Pisarski, Phys. Rev. **D42** (1990) R2156.
- [3] R.D. Pisarski, Phys. Rev. **D47** (1993) 5589. See in the context of QED and with a different approach J.P. Blaizot and E. Iancu, [hep-ph/9706397](#): Phys. Rev. **D55** (1997) 973; Phys. Rev. Lett. 76 (1996) 3080. See also in the context of scalar QED M.H. Thoma and C.T. Traxler, Phys. Lett. **B378** (1996) 233.
- [4] Reference 2; E. Braaten and R.D. Pisarski, Nucl. Phys. **B337** (1990) 569; Phys. Rev. Lett. 64 (1990) 1338; Nucl. Phys. **B339** (1990) 310; J. Frenkel and J.C. Taylor, Nucl. Phys. **B334** (1990) 199; R.D. Pisarski, Phys. Rev. Lett. 63 (1989) 1129; Physica **A** 158 (1989) 246; Nucl. Phys. **A498** (1989) 423c.
- [5] R.D. Pisarski, Physica **A** 158 (1989) 146; Nucl. Phys. **B309** (1988) 476.